

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92560 A2

(51) International Patent Classification⁷: C12Q 1/00

(21) International Application Number: PCT/US01/17554

(22) International Filing Date: 31 May 2001 (31.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/208,405 31 May 2000 (31.05.2000) US

(71) Applicant: **PROMEGA CORPORATION** [US/US];
2800 Woods Hollow Road, Madison, WI 53711-5399
(US).

(72) Inventors: **GOUELI, Said**; 2654 Placid Street, Fitchburg,
WI 53711 (US). **VIDUGIRIENE, Jolanta**; 5719 Long-
ford Terrace, Madison, WI 53711 (US). **KARASSINA,**
Natasha; 2 Sandy Court, Madison, WI 53717 (US).

(74) Agents: **LEONE, Joseph, T. et al.**; DeWitt Ross & Stevens
S.C., 8000 Excelsior Drive, Suite 401, Madison, WI 53717-
1914 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

BEST AVAILABLE COPY

WO 01/92560 A2

(54) Title: ASSAY FOR KINASES AND PHOSPHATASES

(57) Abstract: Disclosed is a method and correspondsind kit for assaying the presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4. The method generally includes the steps of reacting an enzyme with a substrate for a time sufficient to yield phosphorylated or dephosphorylated product; contacting the product with a binding matrix, whereby product is adhered to the matrix; and then analyzing the matrix for presence of, amount of, or both the presence and the amount of the product fixed to the matrix, whereby the presence, the activity, or both the presence and activity of the enzyme can be determined.

ASSAY FOR KINASES AND PHOSPHATASES

PRIORITY

Priority is hereby claimed to provisional application Serial No. 60/208,405, filed 31 May 2000, the entire content of which is incorporated herein.


FIELD OF THE INVENTION

The invention is directed to assays for determining the presence, activity, or both, of kinases and phosphatases. The preferred method is an assay for lipid kinases, phospholipid kinases, and phospholipid phosphatases.

DESCRIPTION OF THE RELATED ART

The importance of phospholipids in general, and phosphoinositides in particular, in the regulation of cellular processes such as cell proliferation, apoptosis, and secretory functions has been recognized for a number of years. While the importance of these compounds is manifest, there is much that remains unknown about how these important cell-signaling compounds are regulated.

Phosphoinositides have the general formula shown in 1:



1

where each R is an unsubstituted or substituted alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group and R^2 , R^3 , R^4 , R^5 and R^6 are hydrogen atoms and their locations on the inositol moiety. These hydrogen atoms can be replaced by phosphate groups singly or in various combinations. The general structure 1 shall be referred to herein as a phosphoinositide, or simply a "PI." Where relevant, the presence and position of phosphate groups on the inositol moiety will be designated by a number designation followed by the letter "P." So, for example, a PI with a phosphate in the R^3 position (*i.e.*, phosphatidylinositol-3-phosphate) will be designated "PI3P," phosphatidylinositol-4-phosphate will be designated "PI4P," etc. For multiply phosphorylated PI's, numbers separated by a comma will be used to designate the positions of the phosphate groups. For example, "PI3,4P₂" designates a PI which is phosphorylated at the R^3 and R^4 positions of the inositol moiety as shown in 1. Phosphorylated PI's in general are referred to as "PIP's"

Because PI's are thought to be central to signal transduction and membrane trafficking in all eukaryotes (Rao et al. (1998) *Cell* 94:829), an understanding of the enzymes that regulate PI's, PIP's, and their metabolites would be extremely helpful. The present invention, an assay to determine the presence and/or activity of lipid kinases, phospholipid kinases, and phosphatases can be used to elucidate further the biological role of PI's.

Any number of assays for measuring enzyme activity are known in the prior art. In particular, Huang et al., U.S. Patent No. 5,869,275, issued February 9, 1999, describes an affinity ultrafiltration-based assay for measuring protein transferase activity. In this approach, labeled and unlabeled substrate having a binding site (the binding site either being created by action of the enzyme being assayed or which exists as an integral part of the substrate, such as an antigenic determinant) are incubated with the enzyme to form labeled product and unlabeled product. The reaction mixture is then contacted with a soluble macroligand capable of forming a specific complex with the product. Of critical importance is that the size of the macroligand-product complex

must be significantly larger than the size of any contaminants or reactants found in the reaction mixture

The macroligand-product complex is then separated from reactants via ultrafiltration. The critical consideration here is that the nominal molecular weight limit of the ultrafiltration membrane must be larger than any potential contaminants in the reaction mixture, as well as larger than any unreacted, labeled substrate, yet smaller than the size of the macroligand-product complex. In this fashion, reactants and contaminants pass through the membrane, while the much larger macroligand-product complex is retained by the membrane. Examination of the ultrafiltration retentate for the presence of labeled product provides an indication of the extent of the reaction.

Mallia, U.S. Patent No. 5,527,688, issued June 19, 1996, describes assays for protein kinases in which a binding membrane is suspended within a reaction vessel, thereby dividing the vessel into two compartments. By adhering products to the suspended membrane, washing can be accomplished centrifugally by placing the wash solution in the upper chamber and centrifuging, thereby forcing the wash solution through the membrane.

SUMMARY OF THE INVENTION

The invention is an assay and a corresponding kit for determining the presence and activity of kinases and phosphatases, and, more specifically, lipid and phospholipid kinases and phosphatases. The preferred embodiment of the invention is an assay capable of measuring the presence and/or activity of any kinase or phosphatase which adds or removes a phosphate group from a lipid or phospholipid substrate.

In particular, a first embodiment of the invention is directed to a method for assaying the presence, the activity, or both the presence and the activity, of an enzyme falling within the enzyme classifications EC 2.7.1, EC 3.1.3, and EC 3.1.4. The method comprises first reacting the enzyme with a corresponding substrate for a time sufficient to yield phosphorylated product when assaying a kinase or a

dephosphorylated product when assaying a phosphatase. The reaction, of course, is run under conditions which render the enzyme under investigation active, and thus the enzyme will catalyze either a phosphorylation (kinase) or a dephosphorylation (phosphatase) of the substrate.

The product formed by the enzymatic reaction is then contacted with a binding matrix. This results in product being bound or fixed to the matrix. With the product fixed on the matrix, the matrix can be mechanically separated from the reaction solution, thereby providing an easy means to separate the products of the enzymatic reaction from unreacted reactant, enzyme, and other non-product ingredients of the reaction solution.

The matrix is then analyzed for the presence of, the amount of, or both the presence and the amount of, the product fixed to the matrix. By determining the presence and/or amount of the product found on the matrix, the presence, the activity, or both the presence and activity of the enzyme that gave rise to the products can be determined.

A second embodiment of the invention is directed to a method as substantially described above, with the addition that the substrate includes a binding moiety. When the substrate is converted into product (by the action of the enzyme), the product also contains the binding moiety.

The product is then contacted with a binding matrix that specifically binds for the binding moiety. This results in product being specifically fixed to the matrix (via the interaction of the binding moiety and the binding matrix. The preferred binding moiety is biotin and the preferred binding matrix is avidin or streptavidin immobilized on an inert support. The matrix is then analyzed as in the first embodiment.

In the second embodiment of the invention, the approach used to modify the lipid or phospholipid enzyme substrate is to include a binding moiety. In this embodiment, the binding moiety (and the binding moiety alone) will bind specifically to a support designed for that purpose. In this fashion, products bearing the binding moiety can be separated easily from products which do not bear the binding moiety by

simply passing the reaction mixture over the support. It is much preferred that the binding moiety be biotin. The support would then comprise any suitable substrate (beads, filter paper, etc.) having avidin or streptavidin immobilized thereto.

In a third embodiment of the invention, a support is not required. In this approach, the assay is conducted entirely in the liquid phase. Using a biotin binding moiety as an example, the avidin or streptavidin would be added to the reaction and the biotin-avidin complexes which form could be separated by any known means (electrophoresis, chromatography, centrifugation, etc.).

In short, the assay of this embodiment functions to determine the presence and activity of lipid and phospholipid kinases as follows: A lipid or phospholipid substrate for the enzyme to be assayed is first modified to include a binding moiety. As noted above, the binding moiety is preferably biotin, although an antibody or antigen can also be used as the binding moiety. Regardless of the choice of binding moiety, it is important that the binding moiety be attached to the substrate in such a fashion that its presence does not interfere with the enzyme's ability to phosphorylate or dephosphorylate the substrate. In most instances, this requires that the binding moiety be added to the end of one of the fatty acyl moieties of the substrate because recognition is dictated largely by the nature of the headgroup.

To assay for lipid and/or phospholipid kinases, the modified substrate is exposed, in the presence of γ - ^{32}P ATP, to a solution thought to contain the kinase of interest and allowed to incubate for a sufficient amount of time and under appropriate conditions such that the kinase, if present and active, can phosphorylate the modified substrate with a ^{32}P -labeled phosphate group. The reaction mixture is then contacted with a capture membrane or matrix, that is, a support bearing a moiety which will capture specifically the modified substrates; in the case of biotin, this would be avidin or streptavidin linked to a support. If binding moiety is an antigen, the capture membrane would include immobilized antibodies specific for the antigen, etc.

When the modified substrates are captured to a solid support, free ^{32}P ATP, reactants, contaminants, etc., are gently washed from the support and the bound

radiolabeled material is measured for radioactivity using a scintillation counter, a "PhosphoImager" device, or by autoradiography.

Where phosphatases are to be assayed, the assay protocol is the same as noted above, with the exception that the substrate is modified to include γ - ^{32}P phosphate groups. In addition, an unlabeled substrate can be used and the released phosphate determined with a colorimetric method or a fluorescent method. (For example, Molecular Probes, Eugene, Oregon, sells a method for fluorescent detection of a released phosphate.) The action of the phosphatase enzyme under analysis will then remove a portion of those groups. In this approach, the reduced activity found on the matrix as compared to the starting labeled substrates is a direct measure of the activity of the phosphatase, or, in the case of non-radiolabeled phosphate, the amount of released phosphate is used as a measure of phosphatase activity.

The invention also comprises a corresponding kit that contains all of the necessary reagents to carry out the assay one or more times. The kit generally comprises: an amount of reaction buffer disposed in a first container; an amount of substrate for an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the substrate disposed in a second container; an amount of purified enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the enzyme disposed in a third container (or the enzyme can be supplied by the user); an amount of binding matrix; and instructions for use of the kit.

A distinct and primary advantage of the assay is that it allows investigators to assay the activity of any number or type of lipid or phospholipid kinases or phosphatases quickly and conveniently.

Another primary advantage of the assay is that it is sufficiently sensitive to assay lipid and phospholipid kinases and phosphatases directly from tissue and cell extracts. Due to the high affinity of the lipid and phospholipid products to the matrix, extraneous free γ ^{32}P -ATP can be removed by washing and the amount of product formed can be determined without need for lipid extraction. This is a distinct

improvement over conventional HPLC and TLC assays, which require lipid extraction. Eliminating lipid extraction, which is costly and time-consuming, makes the subject assay very attractive to investigators in this field.

Another advantage is that the assay is scalable to accommodate high throughput formats. The assay is highly amenable to automation.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1: Flow chart. Parallel flow charts comparing prior art methods to a preferred method according to the present invention.

Fig. 2A: Comparison of PIK activity measurements: present invention vs. conventional phospholipid extraction procedure. Assays were performed using partially-purified PI-3K (500 ng/ml) for 40 minutes or purified PI4P-5K (250 ng/ml) for 20 minutes. For PI-3K activity measurements, PI4,5P₂ (10 µg) + PS (20 µg) was used as reaction substrate. The PI4P-5K reaction was carried out using PI4P (10 µg) as reaction substrate. Samples were processed using the conventional phospholipid extraction procedure illustrate in Fig. 1 (black bars) or according to the present invention (white bars).

Fig. 2B: Comparison of PT-K activity measurements: present invention vs. standard phospholipid extraction procedure. Assays were done using immuno-precipitated PI-3K from liver tissue (500 µg) for 60 min. All the reactions were carried out at room temperature using PIP₂ (10 µg) + PS (20 µg) (lanes 3 and 4) or PI (50 µg) (lanes 1 and 2) as reaction substrates. Samples were processed using the conventional phospholipid extraction procedure illustrated in Fig. 1 (lanes 2 and 4) or according to the present invention (lanes 1 and 3).

Fig. 3: Detection of PI-3K activity associated with activated receptor. 3T3NIH cells (5 x 10⁷ cells) were treated without (lane 1) or with (lane 2) PDGF (50 ng/ml for 5 min at 37°C) and PI-3K was co-immunoprecipitated using anti-phosphotyrosine specific antibodies. The PI-3K reactions were run for 60 min using PIP₂ (10 µg) + PS (20 µg) as substrates.

Figs. 4A and 4B: Dependence of PI-K activity on the amount of lipid substrate loaded on the membrane. **Fig. 4A:** PI-3K activity was measured using PI-3K from Alexis Biochemical (San Diego, California) (12.5 $\mu\text{g/ml}$) for 60 min at room temperature. **Fig. 4B:** PI-5K activity was measured using purified PI-5K (125 ng/ml) for 60 min at room temperature. In addition to lipid substrates, all samples contained 10 μg of PS loaded on the membrane.

Figs. 5A, 5B, 5C, and 5D: Linear detection of PIP-5K activity using the present invention. Assays were performed using the inventive method described herein, utilizing 3 μl of a lipid mixture containing 1 μg of PI4P and 10 μg of PS in chloroform/methanol 2: 1 spotted onto a pre-numbered membrane square. **Figs 5A and 5B:** PI-5K activity was measured at different time points using 4 ng of purified protein. **Figs 5C and 5D:** PI-5K activity was measured for 10 min using different protein concentrations. The reaction products were analyzed by phosphorimaging (**Figs. 5A and 5C**) or by scintillation counting (**Figs 5B and 5D**). Summary: Between 20 pmol to 1.2 nmol of formed product can be detected using the present invention.

Figs. 6A and 6B: Linear detection of PI-3K activity using the present invention. Assays were performed using the inventive method described herein, utilizing 5 μl of PI (25 μg) in chloroform/methanol 2:1 spotted onto a well of a "SAM2"-brand membrane 96-well plate. Assays were performed using decreasing amounts of partially-purified PI-3K: 1 ng, 2 ng, 4 ng, 8 ng, and 16 ng. Based on obtained scintillation counts, the specific activity has been calculated and shown in **Fig 6A**. The calculated initial reaction rate for different amounts of protein is shown in **Fig 6B**.

Figs. 7A, 7B, and 7C: Reproducibility of the subject invention. Assays were performed using: (**Fig. 7A**) partially purified PI-3K (500 ng/ml) and PIP_2 (10 μg) + PS (20 μg) as lipid substrates for 40 minutes; (**Fig. 7B**) PI-5K (125 ng/ml) and PI4P (1 μg) + PS(10 μg) for 15 minutes; (**Fig. 7C**) PI-5K (125 ng/ml) and PI4P (1 μg) for 60 minutes. All assays were performed at room temperature using 1 μCi of ^{32}P -ATP. In **Fig. 7C**, the reactions were carried out on three different plates (samples 1-4; 5-8; 9-12, respectively). In each plate the reactions were done in triplicates at three

different locations. Each point represents the average of three independent reactions. Lighter bars show the average of 9 points done on the same plate.

Fig. 8: Inhibition of PI-3K activity. Reactions according to the present invention were performed using partially-purified PI-3K for different periods of time in the presence (triangles) or absence (diamonds) of the PI-3K inhibitor wortmannin (final concentration 100 nM).

Fig. 9: Reaction specificity toward lipid substrates. Reactions according to the subject invention were performed with purified PI4P-5K (250 ng/ml) for 45 min at room temperature using 1 μ g of different lipid substrates: PI3,4P (lane 1); PI3,5P₂ (lane 2) and PI4,5P₂ (lane 3). In addition to lipid substrates, all samples contained 10 μ g of PS loaded on the membrane.

Fig. 10: Comparison of reaction products: the present invention vs. the conventional phospholipid extraction procedure. Activities were assessed using 1 μ g of PI4P + 10 μ g of PS as reaction substrates and purified PI-5K (2.5 ng) as the enzyme. The reaction was carried out for 10 minutes (lanes 1 and 2) and 0 minutes (lanes 3 and 4). The reaction was worked up using the conventional phospholipid extraction procedure as illustrated in Fig. 1 (lanes 2 and 4) or according to the subject invention (lanes 1 and 3). When the reaction was been performed on membrane sheets, 40% of bound reaction products were re-extracted for TLC analysis.

Fig. 11A: Comparison of PI4P-5 kinase activity using biotinylated and non-biotinylated short-chain lipids and non-biotinylated long-chain lipids. PI4P-5 kinase activity was measured using PI4P-C₆-biotin (lane 1), PI4P-C₈ (lane 2) and PI4P-C₁₆ (lane 3) as reaction substrates. The reaction products were extracted and analyzed via TLC.

Fig. 11B: Comparison of PI-3K activity using biotinylated short-chain lipids and non-biotinylated long-chain lipids. PI-3K activity was measured using PI-C₆-biotin or PI-C₁₆ as reaction substrates. The reaction products were bound to streptavidin-coated membranes. The membranes were washed to remove unreactive reactants and

were analyzed directly (black bars) or were exposed to chloroform:methanol:water (10:10:3) treatment for lipid re-extraction (white bars).

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations & Definitions:

The following abbreviations and definitions are expressly adopted herein. All terms not provided a definition are to be given their accepted definition in the art:

"Alkyl" = a straight, branched, or cyclic, fully-saturated hydrocarbon radical having the number of carbon atoms designated (*i.e.*, C₁-C₂₄ means one to 24 carbons); examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)ethyl, cyclopropylmethyl, and the higher homologs and isomers thereof, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like; an alkyl group will typically contain from 1 to 24 carbon atoms, with those groups having eight or more carbon atoms being preferred in the present invention; a "lower alkyl" is an alkyl group having fewer than eight carbon atoms.

"Alkenyl" = an alkyl group having one or more double bonds or triple bonds; examples of unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butenyl, and the higher homologs and isomers.

"Alkylcarbonyl" = a carbonyl-containing alkyl radical derived from the corresponding carboxylic acid, *e.g.* acyl radicals derived from lauric acid, myristic acid, palmitic acid, stearic acid, and the like; will typically contain from 2 to 24 carbon atoms.

"Alkenylcarbonyl" = a carbonyl-containing alkenyl radical derived from the corresponding carboxylic acid, *e.g.*, acyl radicals derived oleic acid, linoleic acid, linolenic acid, eleostearic acid, arachidonic acid, and the like; will typically contain from 2 to 24 carbon atoms.

"Bn" = benzyl.

"BOM" = benzyloxymethyl.

"EDTA" = ethylenediaminetetraacetic acid

"HEPES" = *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid

"Lipid kinase/phospholipid kinase" = any enzyme falling within the classification EC 2.7.1.x, where "x" is a variable.

"PDGF" = platelet-derived growth factor.

"Phospholipid phosphatase" = any enzyme falling within the classification EC 3.1.3.x and 3.1.4.x, where "x" is a variable.

"PI" = phosphatidylinositol; an unphosphorylated phosphoinositide (*i.e.*, a phosphoinositide lacking any phosphate groups on the inositol moiety).

"PIK" = generally, phosphoinositide kinase; PI-3-kinase, PI-3K, is the illustrative enzyme.

"PIP" = a phosphorylated phosphoinositide (*i.e.*, a PI having one or more phosphate groups present on the inositol moiety); phosphatidylinositol phosphate.

"PI_{x,x',x''}... P_n" = a nomenclature shorthand to designated PIPs, wherein "PI" designates a phosphoinositide, "x, x', x''..." are numerical variables designating the position of phosphate groups on the inositol moiety, "P" designates that the inositol moiety is phosphorylated, and "n" designating the number of phosphate groups present on the inositol moiety.

"PMB" = *p*-methoxybenzyl.

"PS" = phosphatidylserine.

"Substituted" = a radical including one or more substituents, such as lower alkyl, aryl, acyl, halogen, hydroxy, amino, alkoxy, alkylamino, acylamino, thioamido, acyloxy, aryloxy, aryloxyalkyl, mercapto, thia, aza, oxo, both saturated and unsaturated cyclic hydrocarbons, heterocycles and the like; the substituents may be attached to any carbon of the base moiety.

"Substrate" or "corresponding substrate" = a substrate that can be phosphorylated or dephosphorylated by the enzyme being assayed.

"Solid support" = a material that is substantially insoluble in a selected solvent system, or which can be readily separated (*e.g.*, by precipitation) from a selected

solvent system in which it is soluble. Solid supports useful in practicing the present invention can include groups that are activated or capable of activation to allow selected species to be bound to the solid support. A solid support can also be a substrate, for example, a chip, wafer, or well.

"TLC" = thin-layer chromatography.

Modifying Substrates to Contain a Binding Moiety:

Substrates for lipid or phospholipid kinases and phosphatases can be obtained commercially from numerous suppliers.

For example, PI's such as *D-myo*-phosphatidylinositol, *D-myo*-phosphatidylinositol 3-phosphate, *D-myo*-phosphatidylinositol 4-phosphate, *D-myo*-phosphatidylinositol 5-phosphate, *D-myo*-phosphatidylinositol 3,4-bisphosphate, *D-myo*-phosphatidylinositol 3,5-bisphosphate, *D-myo*-phosphatidylinositol 4,5-bisphosphate, *D-myo*-phosphatidylinositol 3,4,5-trisphosphate, and derivatives thereof, PIK antibodies, and biotin-tagged PI's are available commercially from Echelon Research Laboratories Inc., Salt Lake City, Utah; and Upstate Biotechnology, Lake Placid, New York). The PI's noted above, as well as a host of others, derivatives, and antibodies thereto are also available commercially from A.G. Scientific, Inc., San Diego, California.

PI's can also be synthesized using conventional routes from naturally-occurring chiral precursors. For example, inositol head groups can be derived from methyl α -D-glucopyranoside via a Ferrier rearrangement and the diacylglycerol moieties can be prepared from (+)-isopropylideneglycerol.

The most preferred binding moiety to be affixed to the substrate (when such a binding moiety is employed) is a biotin moiety. This is due to biotin's relatively small size, relative ease of chemical manipulation, and its very robust, high-level, and specific binding to avidin and streptavidin.

Biotin can be affixed to an acyl chain of the diacylglycerol portion of a substrate PI via a coupling reaction wherein the biotin is linked via an amide bond to the PI.

Note that when affixed to the acyl chain, the biotin does not interfere with recognition of the modified substrate at both the inositol headgroup and the glycerol backbone proximal to the headgroup. This discovery, that a substrate PI can be modified by the addition of a binding moiety such as biotin without interfering with a lipid/phospholipid kinase's or phosphatase's ability to recognize and bind to the substrate, is novel. In short, no prior art known to the inventors describes or suggests that such a modification can be made.


The preferred reaction to affix biotin (or any other binding moiety having an available reactive group) to a PI is analogous to that described in Chen et al. (1996), 61 *J. Org. Chem.* 6305-6312, incorporated herein by reference. See also G.D. Prestwich (1996), 29 *Acc. Chem. Res.* 503-513, also incorporated herein by reference.

Briefly, lipid-modified analogs of PI's can be formed by inserting an aminoalkanoyl group at the *sn*-1 position of the PI. This group then allows for the addition of a binding moiety, preferably biotin, to the end of the acyl chain.

The synthesis, shown in Reaction Scheme 1, follows a convergent approach, beginning with the selective *sn*-1-*O*-acylation of a protected chiral glycerol synthon, followed by acylation of the *sn*-2 position and oxidative deprotection to yield the desired 1,2-*O*-diacylglycerol derivative.

Reaction with benzyl(*N,N*,*-diisopropylamino*)chlorophosphine yields phosphoramidites, which are then condensed with an appropriately protected *D-myo*-inositol derivative.

REACTION SCHEME 1



In the intermediate compounds, benzyl (Bn) or benzyloxymethyl (BOM) groups protect the final hydroxyl groups and *p*-methoxybenzyl (PMB) groups protect the future phosphomonoesters. Deprotection of the PMB groups, followed by phosphorylation, hydrogenolysis, and ion exchange chromatography yields the aminoacyl modified PI's.

Attachment of the binding moiety via an ester linkage yields a PI having a binding moiety linked thereto.

Where the binding moiety is something other than biotin, such as an antibody or an antigenic determinant, analogous linking chemistries can be utilized to affix the binding moiety to the acyl chain of the PI.

Binding Matrices:

Where no binding moiety is attached to the substrate, the preferred binding matrix is an aldehyde-activated solid support or substrate, most preferably an aldehyde-activated regenerated cellulose. The preferred matrices are "SARTOBIND"®-brand aldehyde membranes, available commercially from Sartorius Corporation (Edgewood, New York, USA and Goettingen, Germany) and SAM2®-brand membranes (described in the following paragraph). Likewise, supports including diethylaminoethyl cellulose and polyvinylidene difluoride can also be used.

Where the binding moiety is biotin, the preferred binding matrix is a support having avidin or streptavidin immobilized thereon. The support may be in the form of a filter, membrane, beads, etc. The most preferred matrices are SAM2® -brand Biotin Capture Membranes or SAM2®-brand 96 Biotin Capture Plates (96-well microtiter format) from Promega Corporation, Madison, Wisconsin. The SAM2®-brand membrane binds biotinylated molecules based on their strong affinity for streptavidin. The process by which the membrane is produced results in a high density of streptavidin on the membrane filter matrix, promoting rapid, quantitative capture of biotinylated substrates. In addition, the SAM2®-brand membrane has been optimized for low nonspecific binding. Using the 96 well plate format allows washes to be performed using a vacuum manifold or a commercially available plate washer.

Where the binding moiety is an antibody or antigenic determinant, the preferred binding matrix is an affinity matrix having immobilized thereon a compound which binds strongly and specifically with the binding moiety.

Enzymes That Can Be Assayed:

The subject assay can be used to detect and measure the presence of any lipid or phospholipid kinase or phosphatase. In short, any enzyme whose catalytic activity transfers a phosphate group to a lipid or phospholipid substrate, or any enzyme whose catalytic activity removes a phosphate group from a lipid or phospholipid substrate, can be assayed using the present invention.

More specifically, the subject assay can be used to determine the presence and/or activity of any lipid or phospholipid kinase falling within the Enzyme Classification (EC) 2.7.1.x (where x is a variable), including, without limitation, EC 2.7.1.1 hexokinase, EC 2.7.1.2 glucokinase, EC 2.7.1.3 ketohexokinase, EC 2.7.1.4 fructokinase, EC 2.7.1.5 rhamnulokinase, EC 2.7.1.6 galactokinase, EC 2.7.1.7 mannokinase, EC 2.7.1.8 glucosamine kinase, EC 2.7.1.10 phosphoglucokinase, EC 2.7.1.11 6-phosphofructokinase, EC 2.7.1.12 gluconokinase, EC 2.7.1.13 dehydrogluconokinase, EC 2.7.1.14 sedoheptulokinase, EC 2.7.1.15 ribokinase, EC 2.7.1.16 ribulokinase, EC 2.7.1.17 xylulokinase, EC 2.7.1.18 phosphoribokinase, EC 2.7.1.19 phosphoribulokinase, EC 2.7.1.20 adenosine kinase, EC 2.7.1.21 thymidine kinase, EC 2.7.1.22 ribosylnicotinamide kinase, EC 2.7.1.23 NAD kinase, EC 2.7.1.24 dephospho-CoA kinase, EC 2.7.1.25 adenylyl-sulfate kinase, EC 2.7.1.26 riboflavin kinase, EC 2.7.1.27 erythritol kinase, EC 2.7.1.28 triokinase, EC 2.7.1.29 glycerone kinase, EC 2.7.1.30 glycerol kinase, EC 2.7.1.31 glycerate kinase, EC 2.7.1.32 choline kinase, EC 2.7.1.33 pantothenate kinase, EC 2.7.1.34 pantetheine kinase, EC 2.7.1.35 pyridoxal kinase, EC 2.7.1.36 mevalonate kinase, EC 2.7.1.37 protein kinase, EC 2.7.1.38 phosphorylase kinase, EC 2.7.1.39 homoserine kinase, EC 2.7.1.40 pyruvate kinase, EC 2.7.1.41 glucose-1-phosphate phosphodismutase, EC 2.7.1.42 riboflavin phosphotransferase, EC 2.7.1.43 glucuronokinase, EC 2.7.1.44 galacturonokinase, EC 2.7.1.45 2-dehydro-3-deoxygluconokinase, EC 2.7.1.46 L-arabinokinase, EC 2.7.1.47 D-ribulokinase, EC 2.7.1.48 uridine kinase, EC 2.7.1.49 hydroxymethylpyrimidine kinase, EC 2.7.1.50 hydroxyethylthiazole kinase, EC 2.7.1.51 L-fuculokinase, EC 2.7.1.52 fucokinase, EC 2.7.1.53 L-xylulokinase,

EC 2.7.1.54 D-arabinokinase, EC 2.7.1.55 allose kinase, EC 2.7.1.56 1-phosphofructokinase, EC 2.7.1.58 2-dehydro-3-deoxygalactonokinase, EC 2.7.1.59 N-acetylglucosamine kinase, EC 2.7.1.60 N-acylmannosamine kinase, EC 2.7.1.61 acyl-phosphate-hexose phosphotransferase, EC 2.7.1.62 phosphoramidate-hexose phosphotransferase, EC 2.7.1.63 polyphosphate-glucose phosphotransferase, EC 2.7.1.64 inositol 1-kinase, EC 2.7.1.65 scyllo-inosamine 4-kinase, EC 2.7.1.66 undecaprenol kinase, EC 2.7.1.67 1-phosphatidylinositol 4-kinase, EC 2.7.1.68 1-phosphatidylinositol-4-phosphate 5-kinase, EC 2.7.1.69 protein-Np-phosphohistidine-sugarphosphotransferase, EC 2.7.1.70 protamine kinase, EC 2.7.1.71 shikimate kinase, EC 2.7.1.72 streptomycin 6-kinase, EC 2.7.1.73 inosine kinase, EC 2.7.1.74 deoxycytidine kinase, EC 2.7.1.75 (now EC 2.7.1.21), EC 2.7.1.76 deoxyadenosine kinase, EC 2.7.1.77 nucleoside phosphotransferase, EC 2.7.1.78 polynucleotide 5'-hydroxyl-kinase, EC 2.7.1.79 diphosphate-glycerol phosphotransferase, EC 2.7.1.80 diphosphate-serine phosphotransferase, EC 2.7.1.81 hydroxylysine kinase, EC 2.7.1.82 ethanolamine kinase, EC 2.7.1.83 pseudouridine kinase, EC 2.7.1.84 alkylglycerone kinase, EC 2.7.1.85 b-glucoside kinase, EC 2.7.1.86 NADH2 kinase, EC 2.7.1.87 streptomycin 3"-kinase, EC 2.7.1.88 dihydrostreptomycin-6-phosphate 3'a-kinase, EC 2.7.1.89 thiamine kinase, EC 2.7.1.90 diphosphate-fructose-6-phosphate 1-phosphotransferase, EC 2.7.1.91 sphinganine kinase, EC 2.7.1.92 5-dehydro-2-deoxygluconokinase, EC 2.7.1.93 alkylglycerol kinase, EC 2.7.1.94 acylglycerol kinase, EC 2.7.1.95 kanamycin kinase, EC 2.7.1.96 (included in EC 2.7.1.86), EC 2.7.1.97 (identical to EC 2.7.1.125), EC 2.7.1.99 {pyruvate dehydrogenase (lipoamide)} kinase, EC 2.7.1.100 5-methylthioribose kinase, EC 2.7.1.101 tagatose kinase, EC 2.7.1.102 hamamelose kinase, EC 2.7.1.103 viomycin kinase, EC 2.7.1.104 diphosphate-protein phosphotransferase, EC 2.7.1.105 6-phosphofructo-2-kinase, EC 2.7.1.106 glucose-1,6-bisphosphate synthase, EC 2.7.1.107 diacylglycerol kinase, EC 2.7.1.108 dolichol kinase, EC 2.7.1.109 {hydroxymethylglutaryl-CoA reductase (NADPH2)} kinase, EC 2.7.1.110 dephospho-{reductase kinase} kinase, EC 2.7.1.111 (now EC

2.7.1.128), EC 2.7.1.112 protein-tyrosine kinase, EC 2.7.1.113 deoxyguanosine kinase, EC 2.7.1.114 AMP-thymidine kinase, EC 2.7.1.115 {3-methyl-2-oxobutanoate dehydrogenase (lipoamide)} kinase, EC 2.7.1.116 {isocitrate dehydrogenase (NADP)} kinase, EC 2.7.1.117 myosin-light-chain kinase, EC 2.7.1.118 ADP-thymidine kinase, EC 2.7.1.119 hygromycin-B kinase, EC 2.7.1.120 caldesmon kinase, EC 2.7.1.121 phosphoenolpyruvate-glycerone phosphotransferase, EC 2.7.1.122 xylitol kinase, EC 2.7.1.123 Ca²⁺/calmodulin-dependent protein kinase, EC 2.7.1.124 {tyrosine 3-monooxygenase} kinase, EC 2.7.1.125 rhodopsin kinase, EC 2.7.1.126 b-adrenergic-receptor kinase, EC 2.7.1.127 1-D-myo-inositol-trisphosphate 3-kinase, EC 2.7.1.128 {acetyl-CoA carboxylase} kinase, EC 2.7.1.129 myosin-heavy-chain kinase, EC 2.7.1.130 tetraacyldisaccharide 4'-kinase, EC 2.7.1.131 low-density-lipoprotein kinase, EC 2.7.1.132 tropomyosin kinase, EC 2.7.1.133 inositol-trisphosphate 6-kinase, EC 2.7.1.134 inositol-tetrakisphosphate 1-kinase, EC 2.7.1.135 tau-protein kinase, EC 2.7.1.136 macrolide 2'-kinase, EC 2.7.1.137 1-phosphatidylinositol 3-kinase, EC 2.7.1.138 ceramide kinase, EC 2.7.1.139 inositol-trisphosphate 5-kinase, EC 2.7.1.140 inositol-tetrakisphosphate 5-kinase, EC 2.7.1.141 {RNA-polymerase}-subunit kinase, EC 2.7.1.142 glycerol-3-phosphate-glucose phosphotransferase, EC 2.7.1.143 diphosphate-purine nucleoside kinase, EC 2.7.1.144 tagatose-6-phosphate kinase, and EC 2.7.1.145 deoxynucleoside kinase.

Preferred kinases which can be assayed using the subject assay include EC 2.7.1.137 (1-phosphatidylinositol 3-kinase, referred to herein as PI-3K), EC 2.7.1.67 (1-phosphatidylinositol 4-kinase), EC 2.7.1.68 (1-phosphatidylinositol-4-phosphate kinase, also called diphosphoinositide kinase or PIP kinase), 1-phosphatidylinositol 5-kinase (PI-5K), and the like.

Regarding phosphatases, the subject assay can be used to determine the presence and/or activity of any lipid or phospholipid phosphatase falling within the Enzyme Classification (EC) 3.1.3.x and EC 3.1.4.x (where x is a variable), including, without limitation, EC 3.1.3.1 alkaline phosphatase, EC 3.1.3.2 acid phosphatase, EC 3.1.3.3

phosphoserine phosphatase, EC 3.1.3.4 phosphatidate phosphatase, EC 3.1.3.5
 5'-nucleotidase, EC 3.1.3.6 3'-nucleotidase, EC 3.1.3.7 3'(2'),5'-bisphosphate
 nucleotidase, EC 3.1.3.8 3-phytase, EC 3.1.3.9 glucose-6-phosphatase, EC 3.1.3.10
 glucose-1-phosphatase, EC 3.1.3.11 fructose-bisphosphatase, EC 3.1.3.12
 trehalose-phosphatase, EC 3.1.3.13 bisphosphoglycerate phosphatase, EC 3.1.3.14
 methylphosphothioglycerate phosphatase, EC 3.1.3.15 histidinol-phosphatase, EC
 3.1.3.16 phosphoprotein phosphatase, EC 3.1.3.17 {phosphorylase} phosphatase, EC
 3.1.3.18 phosphoglycolate phosphatase, EC 3.1.3.19 glycerol-2-phosphatase, EC
 3.1.3.20 phosphoglycerate phosphatase, EC 3.1.3.21 glycerol-1-phosphatase, EC
 3.1.3.22 mannitol-1-phosphatase, EC 3.1.3.23 sugar-phosphatase, EC 3.1.3.24
 sucrose-phosphatase, EC 3.1.3.25 inositol-1(or 4)-monophosphatase, EC 3.1.3.26
 6-phytase, EC 3.1.3.27 phosphatidylglycerophosphatase, EC 3.1.3.28
 ADPphosphoglycerate phosphatase, EC 3.1.3.29 N-acylneuraminate-9-phosphatase,
 EC 3.1.3.30 deleted, included in EC 3.1.3.31, EC 3.1.3.31 nucleotidase, EC 3.1.3.32
 polynucleotide 3'-phosphatase, EC 3.1.3.33 polynucleotide 5'-phosphatase, EC
 3.1.3.34 deoxynucleotide 3'-phosphatase, EC 3.1.3.35 thymidylate 5'-phosphatase,
 EC 3.1.3.36 phosphatidylinositol-bisphosphatase, EC 3.1.3.37
 sedoheptulose-bisphosphatase, EC 3.1.3.38 3-phosphoglycerate phosphatase, EC
 3.1.3.39 streptomycin-6-phosphatase, EC 3.1.3.40
 guanidinodeoxy-scylo-inositol-4-phosphatase, EC 3.1.3.41 4-nitrophenylphosphatase,
 EC 3.1.3.42 {glycogen-synthase-D} phosphatase, EC 3.1.3.43 {pyruvate
 dehydrogenase (lipoamide)}-phosphatase, EC 3.1.3.44 {acetyl-CoA
 carboxylate}-phosphatase, EC 3.1.3.45 3-deoxy-manno-octulosonate-8-phosphatase, EC
 3.1.3.46 fructose-2,6-bisphosphate 2-phosphatase, EC 3.1.3.47
 {hydroxymethylglutaryl-CoA reductase (NADPH)}-phosphatase, EC 3.1.3.48
 protein-tyrosine-phosphatase, EC 3.1.3.49 {pyruvate kinase}-phosphatase, EC
 3.1.3.50 sorbitol-6-phosphatase, EC 3.1.3.51 dolichyl-phosphatase, EC 3.1.3.52
 {3-methyl-2-oxobutanoate dehydrogenase (lipoamide)}-phosphatase, EC 3.1.3.53
 myosin-light-chain-phosphatase, EC 3.1.3.54 fructose-2,6-bisphosphate 6-phosphatase,

EC 3.1.3.55 caldesmon-phosphatase, EC 3.1.3.56 inositol-1,4,5-trisphosphate 5-phosphatase, EC 3.1.3.57 inositol-1,4-bisphosphate 1-phosphatase, EC 3.1.3.58 sugar-terminal-phosphatase, EC 3.1.3.59 alkylacetyl glycerophosphatase, EC 3.1.3.60 phosphoenolpyruvate phosphatase, EC 3.1.3.61 inositol-1,4,5-trisphosphate 1-phosphatase, EC 3.1.3.62 inositol-1,3,4,5-tetrakisphosphate 3-phosphatase, EC 3.1.3.63 2-carboxy-D-arabinitol-1-phosphatase, EC 3.1.3.64 phosphatidylinositol-3-phosphatase, EC 3.1.3.65 inositol-1,3-bisphosphate 3-phosphatase, EC 3.1.3.66 inositol-3,4-bisphosphate 4-phosphatase, EC 3.1.3.67 phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase, EC 3.1.3.68 2-deoxyglucose-6-phosphatase, EC 3.1.4.1 phosphodiesterase I, EC 3.1.4.2 glycerophosphocholine phosphodiesterase, EC 3.1.4.3 phospholipase C, EC 3.1.4.4 phospholipase D, EC 3.1.4.10 1-phosphatidylinositol phosphodiesterase, EC 3.1.4.11 1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase, EC 3.1.4.12 sphingomyelin phosphodiesterase, EC 3.1.4.13 serine-ethanolaminephosphate phosphodiesterase, EC 3.1.4.14 {acyl-carrier-protein} phosphodiesterase, EC 3.1.4.15 adenylyl-{glutamate-ammonia ligase} hydrolase, EC 3.1.4.16 2',3'-cyclic-nucleotide 2'-phosphodiesterase, EC 3.1.4.17 3',5'-cyclic-nucleotide phosphodiesterase, EC 3.1.4.35 3',5'-cyclic-GMP phosphodiesterase, EC 3.1.4.36 1,2-cyclic-inositol-phosphate phosphodiesterase, EC 3.1.4.37 2',3'-cyclic-nucleotide 3'-phosphodiesterase, EC 3.1.4.38 glycerophosphocholine cholinephosphodiesterase, EC 3.1.4.39 alkylglycerophosphoethanolamine phosphodiesterase, EC 3.1.4.40 CMP-N-acylneuramate phosphodiesterase, EC 3.1.4.41 sphingomyelin phosphodiesterase D, EC 3.1.4.42 glycerol-1,2-cyclic-phosphate 2-phosphodiesterase, EC 3.1.4.43 glycerophosphoinositol inositolphosphodiesterase, EC 3.1.4.44 glycerophosphoinositol glycerophosphodiesterase, EC 3.1.4.45 N-acetylglucosamine-1-phosphodiester a-N-acetylglucosaminidase, EC 3.1.4.46 glycerophosphodiester phosphodiesterase, EC 3.1.4.47 variant-surface-glycoprotein phospholipase C, EC 3.1.4.48 dolichylphosphate-glucose phosphodiesterase, EC 3.1.4.49 dolichylphosphate-mannose phosphodiesterase, EC 3.1.4.50 glycoprotein

phospholipase D, EC 3.1.4.51 glucose-1-phospho-D-mannosylglycoprotein phosphodiesterase,

Preferred phosphatases which can be assayed using the subject assay include EC 3.1.3.27 (phosphatidylglycerophosphatase), EC 3.1.3.36 (phosphatidylinositol biphosphatase; triphosphoinositide phosphatase), EC 3.1.3.64 (phosphatidylinositol-3-phosphatase), EC 3.1.3.67 (phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase), EC 3.1.4.10 (1-phosphatidylinositol phosphodiesterase; monophosphatidylinositol phosphodiesterase; phosphatidylinositol phospholipase C), EC 3.1.4.11 (1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase, triphosphoinositide phosphodiesterase), phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase, and the like.

The assay can also be extended to assay for the presence and or activity of phosphorylases.

Kits:

The present invention is also directed to kits that utilize the assay described herein. A basic kit for measuring the presence and/or activity of a lipid/phospholipid kinase or phosphatase enzyme includes a vessel containing a natural, semi-synthetic, or wholly synthetic enzyme substrate and/or a modified enzyme substrate having a binding moiety attached thereto, the modified substrate having specific reactivity to the enzyme to be assayed. The kit also contains a binding matrix that specifically adsorbs or otherwise binds to and immobilizes the binding moiety on the modified substrate. Instructions for use of the kit may also be included. The kit may also include an appropriate amount of reaction buffer disposed in a suitable container.

"Instructions for use," is a tangible expression describing the reagent concentration or at least one assay method parameter such as the relative amount of reagent and sample to be admixed, maintenance time periods for reagent/sample

admixtures, temperature, buffer conditions, and the like. The instructions for use are suitable to enable an analyst to carry out the desired assay.

Where the modified substrate includes the preferred binding moiety, biotin, the preferred embodiment of the kit includes a biotin-binding matrix comprising avidin or streptavidin molecules immobilized on an inert support such as filter disks, beads, plates, or a soluble matrix.

The amounts of the various reagents in the kits can be varied depending on various factors, such as the optimum sensitivity of the assay, the number of assays to be performed, etc.

It is within the scope of the invention to provide manual test kits or test kits for use in automated analyzers.

The Assay Protocol:

Referring now to Fig. 1, which is a flow chart comparing a prior art method (left-hand column) to the preferred embodiment of the present invention (right-hand column), a number of benefits of the present invention are immediately apparent:

Addressing the present invention first, the most immediate advantage is that the lipids do not have to be dried from an organic solvent. Instead, the reaction solution is simply spotted directly onto the binding matrix. The enzyme is added to the bound substrate and the reaction run for a specified period of time. The reaction is then stopped, the membrane is rinsed, and the amount of label retained on the matrix is measured.

In stark contrast, the conventional approach requires that the lipid substrates must be dried from an organic solvent (normally chloroform/methanol), and resuspended (normally by sonication) in an aqueous reaction solution. The enzymatic reaction to be studied is then performed in the aqueous reaction solution. After the reaction is complete, the lipid products must be extracted back into chloroform/methanol for analysis. Thus, the conventional approach entails a number of manual separation steps that are both time-consuming, reagent-consuming, and not

particularly amenable to automation. The need for these drying and extraction steps are minimized or eliminated entirely in the subject invention.

The assay protocol of the present invention is best illustrated via a number of Examples. The Examples are included solely to provide a more complete understanding of the invention described and claimed herein. The Examples do not limit the scope of the claimed invention in any way.

Example 1:

The following Example demonstrates that PI-3 kinase and PI4P-5 kinase, members from two distinct structural families of phosphoinositide kinases (PI and PIP kinases, respectively), can act to add a phosphate to a phospholipid substrate that has been immobilized on a solid support. Moreover, this Example demonstrates that this can be accomplished directly from organic solutions where the reaction substrates exist as monomers (*i.e.*, individual lipid molecules), rather than from an aqueous phase where the reaction substrates exist as vesicles or micelles.

Further, this Example shows that the product lipids remain bound to the matrix during washing procedures, thereby providing an easy means to separate the products of the enzymatic reaction from unreacted reactants, enzyme, and other non-product ingredients of the reaction solution.

The data generated according to the present invention are compared with analogous data generated using the conventional phospholipid extraction procedure exemplified in Fig. 1. In the conventional approach, the substrates for enzyme modification are presented in the form of vesicles or micelles in an aqueous reaction solution. After the enzymatic reaction (which takes place in the aqueous solution), the product lipids are separated from the other reaction components by extracting the products back into an organic phase, normally chloroform/methanol. The product lipids partition into the organic phase, while the other reaction components remain in the aqueous phase. A comparative flow chart of the prior art method and the subject invention is presented in Fig. 1.

Lipids, 10 μg of PI4,5P_2 + 20 μg of PS, were dissolved in chloroform:methanol (2:1) and loaded onto "SAM2"-brand membranes for PI-3K activity measurements. Likewise, 10 μg of PI4P was loaded onto "SAM2"-brand membranes for PI4P-5K activity measurements. The membranes were air dried and a reaction mixture containing 50 mM HEPES/NaOH, pH7.5, 100 mM NaCl, 10 mM MgCl_2 , 20 ng partially-purified PI-3K or 10 ng of purified PI4P-5K, 50 μM ATP supplemented with 1 μCi of $\gamma\text{-}^{32}\text{P}$ -ATP was added onto the membrane containing the immobilized lipids.

Following incubation at room temperature, the reactions were stopped with 7.5 M guanidine chloride and the membranes were washed with 2M NaCl, followed by 2 M NaCl/1 % H_3PO_4 . The washed membranes were dried and subjected to scintillation counting. The obtained data are presented in Fig. 2A (white bars).

In the conventional phospholipid extraction procedure, the same amount of lipids indicated above were dried under nitrogen and re-suspended by sonication in 50 mM HEPES/NaOH, pH 7.5 buffer containing 1 mM EDTA. Then the reaction mixture was added to sonicated lipids and the reaction was carried out in aqueous solution. The reaction was stopped with 1N HCl, lipids were extracted as indicated in Fig. 1 and analyzed. The data are shown in Fig. 2A (black bars).

To generate immunoprecipitated enzyme as used in Fig. 2B, livers were removed from rats after ether anesthesia. The livers were cut into small pieces and homogenized. The homogenate was centrifuged at 10,000 x g for 10 minutes. The supernatant was centrifuged at 15,000 x g for 1 hour. The supernatant was then titrated to pH 5.75 by drop-wise addition of 1 M acetic acid. After stirring for 10 minutes at 4°C, the solution was centrifuged at 10,000 x g for 10 minutes. The pellet was re-suspended in buffer containing 50 mM HEPES/NaOH, pH 7.5.

Then, 6 μl of anti PI-3 kinase p85 rabbit polyclonal IgG (Upstate Biotechnology, Lake Placid, New York) were added to each tube containing the lysed cell solution and the tubes were incubated overnight at 4°C. Then 100 μl of protein A sepharose CL-4B (Pharmacia, Peapack, New Jersey) was added to each tube and they

were further incubated for 2 hours at 4°C. The sepharose/antibody/antigen complex (the "complex") was then pelleted by centrifugation, the supernatant removed and the complex washed twice with PBS containing 1 % NP-40 and 10 % glycerol, three times with 100mM Tris/HCl (pH 7.5) containing 500 mM NaCl and 100 μ M Na_3VO_4 and twice with 10mM Tris/HCl containing 100 mM NaCl, 1 mM EDTA, 100 μ M Na_3VO_4 . Then 50 μ l of 10 mM Tris/HCl (pH 7.5) containing 100 mM NaCl was added to the complex and this solution was then referred to as the immunoprecipitated PI-3 kinase enzyme..

Immunoprecipitated PI-3 K was assayed as described above for purified PI-3K with lipid substrates immobilized on "SAM"-brand membranes. All reactions were carried out at room temperature using 10 μ g PI4,5P2 + 20 μ g PS (Fig. 2B, bars 3 and 4) or 50 μ g PI (Fig. 2B, bars 1 and 2) as reaction substrates.

Example 2:

An assays analogous to that depicted in Example 1 can be assembled to assay for the presence and/or activity of lipid and phospholipid phosphatases, such as phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase.

Here, the substrate bearing the binding moiety is phosphorylated with a known amount of $\{\gamma\text{-}^{32}\text{P}\}$ ATP.

The reaction components were the same as described in Example 1. The reaction components were applied directly to a "SAM2"-brand membrane and the reaction run for the specified period of time and then terminated. The results of the experiment are generated by measuring the decreased amount of radiation present in the products which adhere to the binding matrix as compared to the radioactivity present in the reactants.

Example 3:

In resting cells, PI-3K is located in the cytosol. However, upon cell stimulation, the enzyme is recruited to the plasma membrane where it is associated with

particular receptors and is involved in further propagation of the signal. This example illustrates that the described method allows to measure PI-3K activity associated with activated receptor. 3T3 NIH cells (0.5×10^7 cells) were starved overnight in serum-free medium. PDGF (50 ng/ml) was added to the starved cells and the induction was carried out for 5 min at 37°C. Following the PDGF induction, cells were washed once with PBS while on the plate, scraped into PBS, transferred to a centrifuge tube, pelleted by centrifugation. The cell pellet was lysed and activated receptor-PI-3K complex was immunoprecipitated using anti-phosphotyrosine specific antibodies. The PI-3K reaction was run for 60 min using PI4,5P₂ (10 µg) as substrate, immobilized on "SAM2"-brand membranes, together with 20 µg of carrier lipid (PS). The increase in PI-3K activity upon stimulation with PDGF is shown in Fig. 3. This Example clearly indicates that the present invention enables cells to be monitored for activation and also enables the analysis of PI-3K activity associated with activated receptors.

Example 4:

This Example illustrates the dependence of PIK activity on the amount of lipid substrate loaded onto the binding matrix. In Fig. 4A, PI-3K activity was measured using PI-3K from Alexis (12.5 µg/ml) for 60 min at room temperature (using the protocol of Example 1, with any modifications noted). In Fig. 4B, PI-5K activity was measured using purified PI-5K (125 ng/ml) for 60 min at room temperature. In addition to lipid substrates, all samples contained 10 µg of PS loaded on the membrane.

As shown in Fig. 4A, the reaction yields a linear response from 0 to roughly 6 µg of substrate. The reaction then reaches a saturation point at roughly 7 µg of PIP₂. Beyond this concentration of substrate, the presence of additional substrate does not result in higher radioactivity incorporation in the formed product. Fig. 4B, which illustrates the results for PI-5K, shows that the enzyme activity expressed as CPM count remains unchanged in going from PI4P concentrations from 1 to 10 µg, thus indicating that at these substrate levels, the reaction is already saturated.

Example 5:

This Example illustrates the linear detection of PIP-5K activity using the present invention. Assays were performed using the described protocol (Example 1) with 3 μ l of lipid mixture containing 1 μ g of PI4P and 10 μ g of PS in chloroform/methanol 2:1 spotted onto a pre-numbered membrane square.

In Figs. 5A and 5B, the PI-5K activity was measured at different time points (0, 2.5, 5, 10, and 20 min) using 4 ng of purified protein. In Figs. 5C and 5D, PI-5K activity was measured for 10 min using different protein concentrations (0, 50, 125, 200, 250 ng/ml). The reaction products were analyzed by phosphorimaging (Figs. 5A and 5C) or by scintillation counting (Figs 5B and 5D). As is clearly illustrated by this series of figures, the assay protocol yields linear time-dependent results (Figs 5A and 5B) and concentration-dependent results (Figs. 5C and 5D). Between 20 pmol and 1.2 nmol of formed product can be detected using the subject invention.

Example 6:

In this Example, linear detection of PI-3K activity was assayed using the present invention. Assays were performed using the described protocol with 5 μ l of PI (25 μ g) in chloroform/methanol 2:1 spotted onto a well of modified SAM-brand membrane 96-well plate (Promega). Assays were performed using different amounts of partially purified PI-3K: 1 ng (light green); 2 ng (dark green); 4 ng (yellow); 8 ng (orange); 16 ng (red). Based on obtained scintillation counts, the specific activity has been calculated and shown in Fig. 6A). The calculated initial reaction rate for different amount of protein is shown in Fig. 6B. This Example shows that the subject method yields linear detection of enzyme over a broad concentration range.

Example 7:

This Example is a comparison of reaction products detected using the present invention versus the conventional phospholipid extraction procedure outlined in Fig. 1. In Fig. 10, activities were assessed using 1 μ g of PI4P + 10 μ g of PS as reaction substrates and purified PI-5K as an enzyme source. The reaction was carried out for 0 (lanes 3, 4) and 30 minutes (lanes 1, 2). The reaction was performed using a conventional phospholipid extraction procedure (lanes 2, 4) or according to the subject invention (lanes 1, 3). When using the subject invention, following reaction performed on membrane sheets, the lipids were re-extracted for TLC analysis.

The results show that the products formed using the present invention yield useful information on enzymatic activity, information that is comparable to the widely-used conventional assay described earlier.

Example 8:

This Example demonstrates the reproducibility of the subject invention. Assays were performed using:

Fig. 7A: partially purified PI-3K (500 ng/ml) and PIP₂ (10 μ g) + PS (20 μ g) as lipid substrates for 40 minutes

Fig. 7B: PI-5K (125 ng/ml) and PI4P (1 μ g) + PS(10 μ g) for 15 minutes

Fig. 7C: PI-5K (125 ng/ml) and PI4P (1 μ g) for 60 minutes.

All assays were performed at room temperature using 1 μ Ci of ³²P-ATP.

In Fig. 7C, the reactions were carried out on three different plates (samples 1-4; 5-8; 9-12, respectively). In each plate the reactions were done in triplicates at three different locations. Each point represents the average of three independent reactions. Yellow bars show the average of 9 points done on the same plate.

As is clearly shown by this set of experiments, the method is highly reproducible.

Example 9:

This Example illustrates that the products formed by the enzymatic reaction are due solely to the presence of active enzyme. This is shown by the loss of products formed when the reaction is run in the presence of an inhibitor that is specific to PI-3K.. A reaction was run with partially purified PI-3K for different periods of time in the presence (triangles) or absence (diamonds) of PI-3K inhibitor wortmannin (final concentration 100 nM). See Fig. 8. As shown in Fig. 8, the present method yielded greatly reduced radiolabeled product as expressed in CPM values when the assay was performed in the presence of the PI-3K inhibitor.

Example 10:

This Example illustrates the reaction specificity toward lipid substrates. Here, a reaction was performed with purified PI-5K (125 ng/ml) for 15 min at room temperature using different lipid substrates: PI3,4P (lane 1); PI3,5P (lane 2) and PI4,5P2 (Lane 3). PI3,5P (lane 2) and PI4,5P (lane 3), which already have a phosphate group in the 5-position, are not substrates for PI-5K, an enzyme that functions to catalyze the addition of a phosphate group to the 5-position of a suitable substrate. Thus, as clearly shown in Fig. 9, the assay generate appropriate results and thus demonstrates specificity for enzyme substrates. PI3,4P (lane 1) is a suitable substrate for the PI-5K enzyme and shows the expected, corresponding CPM values, indicating that the enzyme has, in fact, specifically phosphorylated these substrates.

Example 11:

The following Example demonstrates that PI-3 kinase (PI-3K) can act to add a phosphate to a biotinylated phospholipid when the phospholipid is biotinylated in such a way as not to interfere with the interaction between the enzyme and the biotinylated substrate. The biotin group is attached to the lipid substrate, preferably to the end of the lipid chain. This Example also demonstrates that a streptavidin matrix is capable of separating the biotinylated, phosphorylated product from non-biotinylated

components of the reaction mixture to allow for detection and quantitation of the product.

Lipids: 30 μ g of short-chain PI4P-C₈, biotin-modified short-chain PI4P-C₆-biotin, and long-chain PI4P-C₁₆ were dried under vacuum in the presence of 10 μ g carrier lipid (PS) in separate tubes. The dried lipids were dissolved in 50 mM HEPES/NaOH, pH 7.5 + 1 mM EDTA buffer and subjected to sonication for 5 min. This solution is referred to as the "prepared substrate."

The following kinase reactions were then assembled:

4 μ l of 100 mM MgCl₂

20 μ l of prepared substrate

1 μ l of purified PI4P-5K or PI-3K

11 μ l 50 mM HEPES/NaOH + 100 mM NaCl buffer

4 μ l of 0.5 mM ATP containing 1 μ Ci ³²P-ATP

The reactions were carried out for 15 minutes at room temperature. In Fig. 11A, following the reaction, lipids were extracted and analyzed on TLC to show that biotinylated lipids can be modified by PI-kinases. In Fig. 11B, 25 % of the reaction mixture was loaded on "SAM2"-brand membranes, the membranes were washed and then analyzed directly (black bars) or the membranes were exposed for additional treatment with chloroform/methanol/water (10:10:3) to remove and separate modified from unmodified lipids (white bars).


This Example indicates that membrane-bound lipids remain attached to the matrix during separation procedures and thereby allows separation of reaction products from other reaction components. However, natural lipids bound to the membranes can be removed under defined conditions, for example, following membrane treatment with chloroform:methanol:water (10:10:3). Thus, the reaction products can be subjected to more detailed structural analysis if needed.

CLAIMS

What is claimed is:

1. A method for assaying presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the method comprising:
 - (a) under conditions which render the enzyme active, reacting the enzyme with a corresponding substrate for a time sufficient to yield phosphorylated product when assaying a kinase or a dephosphorylated product when assaying a phosphatase; and
 - (b) contacting the product with a binding matrix; whereby product is fixed to the matrix; and then
 - (c) analyzing the matrix for presence of, amount of, or both the presence and the amount of the product fixed to the matrix, whereby the presence, the activity, or both the presence and activity of the enzyme can be determined.
2. The method of Claim 1, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 2.7.1.67, EC 2.7.1.68, and EC 2.7.1.137.
3. The method of Claim 2, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated support.
4. The method of Claim 2, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated regenerated cellulose support.


5. The method of Claim 2, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
6. The method of Claim 5, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of ^{32}P -labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phospho-imager.
7. The method of Claim 2, wherein in step (a), the enzyme is reacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C_2 to C_{24} alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and R^6 are selected from the group consisting of hydrogen and phosphate, provided that not all of $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and R^6 are simultaneously phosphate.

8. The method of Claim 1, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 3.1.3.27, EC 3.1.3.36, EC 3.1.3.64, EC 3.1.3.67, EC 3.1.4.10, and EC 3.1.4.11.
9. The method of Claim 8, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated support.


10. The method of Claim 8, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated regenerated cellulose support.
11. The method of Claim 8, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
12. The method of Claim 11, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of ^{32}P -labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phospho-imager.
13. The method of Claim 8, wherein in step (a), the enzyme is reacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C_2 to C_{24} alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and R^6 are selected from the group consisting of hydrogen and phosphate, provided that not all of $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and R^6 are simultaneously hydrogen.

14. A method for assaying presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the method comprising:
- (a) under conditions which render the enzyme active, reacting the enzyme with a corresponding substrate, the substrate including a binding moiety, for a time sufficient to yield phosphorylated product when assaying a kinase or a dephosphorylated product when assaying a phosphatase, the product also including the binding moiety; and
 - (b) contacting the product with a binding matrix specific for the binding moiety; whereby product is specifically fixed to the matrix; and then
 - (c) analyzing the matrix for presence of, amount of, or both the presence and the amount of the product, whereby the presence, the activity, or both the presence and activity of the enzyme can be determined.
15. The method of Claim 14, wherein the binding moiety is biotin and the binding matrix is avidin or streptavidin immobilized on a solid support.
16. The method of Claim 14, wherein the binding moiety is an antigenic determinant and the binding matrix is an antibody specific for the antigenic determinant, the antibody being immobilized on a solid support.
17. The method of Claim 14, wherein the binding moiety is an antibody, and the binding matrix is an antigenic determinant specific to the antibody, the antigenic determinant being immobilized on a solid support.
18. The method of Claim 14, wherein the binding moiety is an antibody, and the binding matrix is an anti-antibody specific for the antibody, the anti-antibody being immobilized on a solid support.


19. The method of Claim 14, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 2.7.1.67, EC 2.7.1.68, and EC 2.7.1.137.
20. The method of Claim 19, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated support.
21. The method of Claim 19, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated regenerated cellulose support.
22. The method of Claim 19, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
23. The method of Claim 22, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of ^{32}P -labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phospho-imager.
24. The method of Claim 19, wherein in step (a), the enzyme is reacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C₂ to C₂₄ alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and R², R³, R⁴, R⁵ and R⁶ are selected from the group consisting of hydrogen and phosphate, provided that not all of R², R³, R⁴, R⁵ and R⁶ are simultaneously phosphate.

25. The method of Claim 14, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 3.1.3.27, EC 3.1.3.36, EC 3.1.3.64, EC 3.1.3.67, EC 3.1.4.10, and EC 3.1.4.11.
26. The method of Claim 25, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated support.
27. The method of Claim 25, wherein in step (b), the product is contacted with a binding matrix comprising an aldehyde-activated regenerated cellulose support.
28. The method of Claim 25, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
29. The method of Claim 28, wherein in step (a), the enzyme is reacted with the corresponding substrate in the presence of ³²P-labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phospho-imager.


30. The method of Claim 25, wherein in step (a), the enzyme is reacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C₂ to C₂₄ alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and R², R³, R⁴, R⁵ and R⁶ are selected from the group consisting of hydrogen and phosphate, provided that not all of R², R³, R⁴, R⁵ and R⁶ are simultaneously hydrogen.

31. A method for assaying presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the method comprising:
- (a) contacting an enzyme substrate specifically reactive with an enzyme of a classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4 with a binding matrix; whereby the enzyme substrate is fixed to the matrix; and then
 - (b) contacting the substrate fixed to the matrix with an enzyme under conditions wherein the enzyme is active for a time sufficient to yield phosphorylated product fixed to the matrix when assaying a kinase or a dephosphorylated product fixed to the matrix when assaying a phosphatase; and then
 - (c) analyzing the matrix for presence of, amount of, or both the presence and the amount of the product fixed to the matrix, whereby the presence, the activity, or both the presence and activity of the enzyme can be determined.


32. The method of Claim 31, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 2.7.1.67, EC 2.7.1.68, and EC 2.7.1.137.
33. The method of Claim 32, wherein in step (a), the enzyme substrate is contacted with a binding matrix comprising an aldehyde-activated support.
34. The method of Claim 32, wherein in step (a), the enzyme is contacted with a binding matrix comprising an aldehyde-activated regenerated cellulose support.
35. The method of Claim 32, wherein in step (b), the substrate is contacted with the enzyme in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
36. The method of Claim 35, wherein in step (b), the substrate is contacted with the enzyme in the presence of ^{32}P -labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phosphorimager.
37. The method of Claim 32, wherein in step (a), the matrix is contacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C_2 to C_{24} alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and

R^6 are selected from the group consisting of hydrogen and phosphate, provided that not all of R^2, R^3, R^4, R^5 and R^6 are simultaneously phosphate.


38. The method of Claim 31, wherein the enzyme assayed is classified within an enzyme classification selected from the group consisting of EC 3.1.3.27, EC 3.1.3.36, EC 3.1.3.64, EC 3.1.3.67, EC 3.1.4.10, and EC 3.1.4.11.
39. The method of Claim 38, wherein in step (a), the substrate is contacted to a binding matrix comprising an aldehyde-activated support.
40. The method of Claim 38, wherein in step (a), the substrate is contacted to a binding matrix comprising an aldehyde-activated regenerated cellulose support.
41. The method of Claim 38, wherein in step (b), the substrate is contacted with the enzyme in the presence of labeled phosphate groups, and in step (c), the matrix is analyzed by determining the presence, the amount, or the presence and amount, of labeled phosphate groups fixed to the matrix.
42. The method of Claim 41, wherein in step (b), the substrate is contacted with the enzyme in the presence of ^{32}P -labeled phosphate groups, and in step (c), the matrix is analyzed using a scintillation counter or a phospho-imager.
43. The method of Claim 38, wherein in step (a), the matrix is contacted with a substrate of formula:



wherein each R is independently an unsubstituted or substituted C₂ to C₂₄ alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and R², R³, R⁴, R⁵ and R⁶ are selected from the group consisting of hydrogen and phosphate, provided that not all of R², R³, R⁴, R⁵ and R⁶ are simultaneously hydrogen.

44. The method of Claim 31, wherein in step (a), the enzyme substrate is contained within a cell lysate and the cell lysate is contacted with the matrix.
45. The method of Claim 31, wherein in step (a), the enzyme substrate is contained within an organic-phase solution and the organic-phase solution is contacted with the matrix.
46. The method of Claim 31, wherein in step (a), the enzyme substrate is contacted with the matrix in the absence of drying the substrate and in the absence of extracting the substrate from an organic phase into an aqueous phase prior to contacting it with the matrix.
47. A kit for assaying presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the kit comprising:
 - an amount of reaction buffer disposed in a first container;
 - an amount of substrate for an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the substrate disposed in a second container;
 - an amount of binding matrix; and
 - instructions for use of the kit.

48. The kit of Claim 47, wherein the substrate is of formula:



wherein each R is independently an unsubstituted or substituted C₂ to C₂₄ alkyl, alkenyl, alkylcarbonyl, or alkenylcarbonyl group, and R², R³, R⁴, R⁵ and R⁶ are selected from the group consisting of hydrogen and phosphate.

49. The kit of Claim 47, wherein the binding matrix is aldehyde-activated regenerated cellulose.
50. The kit of Claim 47, wherein the substrate further comprises a binding moiety attached thereto.
51. The kit of Claim 50, wherein the binding moiety is biotin.
52. The kit of Claim 50, wherein the binding matrix is avidin or streptavidin immobilized on an inert support.
53. The kit of Claim 47, further comprising an amount of purified enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4, the enzyme disposed in a third container;

1/11

Standard Extraction Method
PRIOR ART

Dry lipids under nitrogen
↓
Sonicate in buffer to form micelles
Add reaction mixture
↓
Stop Reaction
↓
Extract lipids in chloroform/methanol
to induce two phases
↓
Spin down
↓
Remove upper (water) phase
↓
To lower phase add
artificial upper phase
↓
Spin down
↓
Remove lower (lipid) phase
into fresh tubes
↓
Count or run TLC

Present Invention

Spot liquids on the membrane directly
from chloroform/methanol solution
↓
Add reaction mixture
(enzyme + ^{32}P -ATP)
↓
Stop reaction
↓
Wash membranes
↓
Phosphoimage analysis or
radioactivity counting

FIG. 1

2/11




FIG. 2A




FIG. 2B

3/11




FIG. 3

4/11




FIG. 4A




FIG. 4B

5/11




FIG. 5A




FIG. 5C




FIG. 5B




FIG. 5D

6/11




FIG. 6A




FIG. 6B

7/11




FIG. 7A




FIG. 7B




FIG. 7C

8/11




FIG. 8

9/11




FIG. 9




FIG. 10

11/11




FIG. 11A




FIG. 11B

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/092560 A3

- (51) International Patent Classification⁷: C12Q 1/00, 1/48, 1/42, 1/44 (72) Inventors: GOUELI, Said; 2654 Placid Street, Fitchburg, WI 53711 (US). VIDUGIRIENE, Jolanta; 5719 Longford Terrace, Madison, WI 53711 (US). KARASSINA, Natasha; 2 Sandy Court, Madison, WI 53717 (US).
- (21) International Application Number: PCT/US01/17554
- (22) International Filing Date: 31 May 2001 (31.05.2001) (74) Agents: LEONE, Joseph, T. et al.; DeWitt Ross & Stevens S.C., 8000 Excelsior Drive, Suite 401, Madison, WI 53717-1914 (US).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/208,405 31 May 2000 (31.05.2000) US (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (71) Applicant: PROMEGA CORPORATION [US/US]; 2800 Woods Hollow Road, Madison, WI 53711-5399 (US). (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

[Continued on next page]

(54) Title: ASSAY FOR KINASES AND PHOSPHATASES

Standard Extraction Method
PRIOR ART

Dry lipids under nitrogen
↓
Sonicate in buffer to form micelles
Add reaction mixture
↓
Stop Reaction
↓
Extract lipids in chloroform/methanol
to induce two phases
↓
Spin down
↓
Remove upper (water) phase
↓
To lower phase add
artificial upper phase
↓
Spin down
↓
Remove lower (lipid) phase
into fresh tubes
↓
Count or run TLC

Present Invention

Spot liquids on the membrane directly
from chloroform/methanol solution
↓
Add reaction mixture
(enzyme + ³²P-ATP)
↓
Stop reaction
↓
Wash membranes
↓
Phosphoimage analysis or
radioactivity counting

(57) Abstract: Disclosed is a method and corresponding kit for assaying the presence, activity, or both, of an enzyme classified within an enzyme classification selected from the group consisting of EC 2.7.1, EC 3.1.3, and EC 3.1.4. The method generally includes the steps of reacting an enzyme with a substrate for a time sufficient to yield phosphorylated or dephosphorylated product; contacting the product with a binding matrix, whereby product is adhered to the matrix; and then analyzing the matrix for presence of, amount of, or both the presence and the amount of the product fixed to the matrix, whereby the presence, the activity, or both the presence and activity of the enzyme can be determined.

WO 01/092560 A3



patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:

1 August 2002

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/17554

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/00 C12Q1/48 C12Q1/42 C12Q1/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, MEDLINE, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 00584 A (UNIVERSITY OF UTAH RESEARCH FOUNDATION) 6 January 2000 (2000-01-06) page 4, line 8 -page 6, line 13 page 12, line 28 -page 14, line 26 the whole document	31,32, 35,36, 44-46
Y	---	1-53
X	EP 0 444 302 A (BECTON DICKINSON AND COMPANY) 15 May 1996 (1996-05-15) page 4, line 6 -page 4, line 46 ---	1,47,53
X	US 5 869 275 A (HUANG) 9 February 1999 (1999-02-09) column 3, line 50 -column 3, line 64 --- -/--	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

22 May 2002

Date of mailing of the international search report

13/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Thumb, W

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/17554

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 23612 A (PROMEGA CORPORATION) 8 September 1995 (1995-09-08) page 4, line 19 -page 5, line 18 page 6, line 14 -page 7, line 32 page 16, line 17 -page 19, line 16 ----	1-53
Y	DA-SHENG WANG ET AL.: "Biotinylated phosphatidylinositol 3,4,5-triphosphate as affinity ligand" ANALYTICAL BIOCHEMISTRY, vol. 280, 1 May 2000 (2000-05-01), pages 301-307, XP002198359 the whole document ----	1-53
Y	CHAUDHARY, A. ET AL.: "Rapid purification of reporter group-tagged inositol hexakisphosphate on ion-exchange membrane absorbers" BIOTECHNIQUES, vol. 23, September 1997 (1997-09), pages 427-430, XP001071257 the whole document ----	3,4,9, 10,20, 21,26, 27,33, 34,39, 40,49
A	ERNEUX, C. ET AL., : "The diversity and possible functions of the inositol polyphosphate 5-phosphatases" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1436, 1998, pages 185-199, XP004277225 the whole document ----	1,8-18, 25-31, 38-53
A	MAEHAMA, T., AND DIXON, J.E.: "The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-triphosphate" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 22, 29 May 1998 (1998-05-29), pages 13375-13378, XP002088259 the whole document ----	1-53
A	WO 97 40173 A (CHIRON CORPORATION) 30 October 1997 (1997-10-30) the whole document ----	1-53
A	DE 199 42 268 A (HANS-KNÖLL-INSTITUT FÜR NATURSTOFF-FORSCHUNG) 30 March 2000 (2000-03-30) the whole document -----	1-53

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/17554

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0000584	A	06-01-2000	AU 4724099 A WO 0000584 A2	17-01-2000 06-01-2000
EP 0444302	A	04-09-1991	US 5279935 A AT 138201 T AU 625790 B2 AU 6701790 A CA 2030649 A1 DE 69027020 D1 DE 69027020 T2 DK 444302 T3 EP 0444302 A1 ES 2087118 T3 FI 911013 A GR 3020247 T3 IE 904127 A1 JP 4249768 A JP 6105254 B	18-01-1994 15-06-1996 16-07-1992 05-09-1991 02-09-1991 20-06-1996 31-10-1996 02-09-1996 04-09-1991 16-07-1996 02-09-1991 30-09-1996 11-09-1991 04-09-1992 21-12-1994
US 5869275	A	09-02-1999	NONE	
WO 9523612	A	08-09-1995	US 6348310 B1 AU 2097595 A EP 0760678 A1 JP 10511260 T WO 9523612 A1 US 6066462 A	19-02-2002 18-09-1995 12-03-1997 04-11-1998 08-09-1995 23-05-2000
WO 9740173	A	30-10-1997	AU 2438197 A EP 0909325 A2 WO 9740173 A2	12-11-1997 21-04-1999 30-10-1997
DE 19942268	A	30-03-2000	DE 19942268 A1	30-03-2000

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)